

ANA VEIGA

580010683

### ROLE IN THE PROJECT

Review and examine all published data of seasonality pattern and duration across four species of sea turtles, namely, loggerhead (*Caretta caretta*), leatherback (*Dermochelys coriacea*), hawksbill (*Eretmochelys imbricata*) and green turtle (*Chelonia mydas*) at a global scale. Analyse the data and do the write-up.

### PROBLEMS EXPERIENCED

Originally this project was designed to include all the seven species of sea turtles. However, due to lack of reliable published data from the restricted number of nesting sites under study for olive ridley (*Lepidochelys olivacea*), Kemp's ridley (*Lepidochelys kempi*) and flatback turtle (*Natator depressus*), these species were excluded from this study.

There were also important nesting sites that were not included in this study because seasonality of nesting data was not available or the available data did not cover the full nesting season.

### TARGET JOURNAL

Marine Ecology Progress Series

### REASONS FOR CHOICE OF THE TARGET JOURNAL

This happens to be one of the leading journals in the field of marine ecology. It maintains a high standard of quality for scientific contributions and contains relevant topical research studies. I have chosen this journal because I favour its writing style and I feel the topics it publishes are closest to my topic of research.

# Seasonality and Temperature: Turtles in Hot Water?

Ana Veiga<sup>1</sup>

<sup>1</sup> Centre for Ecology and Conservation, School of Biosciences, University of Exeter, Cornwall Campus, Penryn TR10 9EZ, UK

Email: amv202@exeter.ac.uk

**ABSTRACT:** Variation in temperature affects the biology of sea turtles at a range of scales. To elucidate the drivers of seasonality of nesting and duration of season, databases across four species of sea turtles (*Caretta caretta* n=37, *Chelonia mydas* n=64, *Dermochelys coriacea* n=44 and *Eretmochelys imbricata* n=36) at a global scale were created. By using remotely sensed sea surface temperature data, thermal profiles across the nesting season were generated. Duration of nesting season was correlated with latitude in all species but was more tightly coupled with temperature; seasons were significantly longer with increased mean SST. In general, nesting seasonality occurred at warmest time of the year. SST for the month before, month after and the month of peak nesting significantly affected the month of peak nesting.

**KEY WORDS:** Seasonality • Sea Surface Temperature • Latitude • *Caretta caretta* • *Chelonia mydas* • *Dermochelys coriacea* • *Eretmochelys imbricata* • Sea turtles

## INTRODUCTION

As with all ectothermic animals, temperature has a profound influence on the life history of sea turtles (Davenport 1997). Variation in temperature affects the biology of this group at a range of scales. Sea turtles are globally distributed throughout the tropical and subtropical seas (Mazaris et al. 2008) and may experience and tolerate widely different temperatures (Mrosovsky 1980). For instance, leatherbacks (*Dermochelys coriacea*) swim far into the northern and southern hemispheres, loggerhead turtles (*Caretta caretta*) have a sub-tropical nesting range (Mrosovsky 1980), Kemp's ridley (*Lepidochelys kempii*) and other species are found in New England waters (Lazell, 1980). Adult sea turtles migrate hundreds to thousands of kilometres between specific feeding and breeding areas (Plotkin 2003). In addition, at the start of cold season, they migrate toward warmer waters and to avoid cold stunning, larger turtles may enter a hibernation-like state (Milton & Lutz 2003). It has also been reported that during the overwintering phase, there is an increase in the period spent on the seabed for green turtles and loggerheads as a result of long resting dives (Broderick et al. 2007). Moreover, hyperthermia is unusual when turtles are in the ocean but can be experienced while they are on land, basking or nesting (Milton & Lutz 2003). Thus, both high and low

temperatures have negative impacts on sea turtles physiology, affecting feeding behaviour, acid-base and ion balance, and stress hormone levels (Milton & Lutz 2003). Food availability and temperature influence sea turtles' growth rates (Heppell et al. 2003) which is decelerated by cold temperature (Mrosovsky 1980). Temperature also has a profound impact on their reproduction in several ways which I outline in turn.

Sea turtles show temperature-dependent sex determination (TSD) (Bull 1980), with cooler nests producing predominantly males and warm nests producing predominantly females (Morreale et al. 1982). Approximately equal numbers of males and females are produced at pivotal temperatures (Yntema & Mrosovsky 1982). Furthermore, the sex ratio of offspring differs at different times of the year (Mrosovsky et al. 1984). The period when gender is determined by temperature occurs during the middle third of development of embryos (Standora & Spotila 1985). Temperature also influences the duration of incubation (Miller 1985), it is longer at cooler temperatures (Mrosovsky 1980).

In general, female sea turtles do not breed in consecutive years (Davenport 1997), but spend two or more years feeding at sea between successive nesting years (Solow et al. 2002). Males and females do not necessarily have to use the same time scale for reproduction (Miller 1997). The time for two successive seasons at the beach is defined as remigration interval (Carr & Carr 1972). Moreover, the mean remigration interval reported for females differs among the species (Miller 1997). Solow et al. (2002) reported that increased sea surface temperature (SST) has a positive effect on the 2-year remigration probability in the second winter following nesting for green turtles (*Chelonia mydas*) in Tortuguero, Costa Rica, with thermally dependent process affecting nutritional status in the previous winter sufficient to sustain vitellogenesis. For Pacific loggerhead population, Chaloupka et al. (2008) reported that there was a significant inverse correlation between nesting abundance and mean SST, because warmer foraging habitats are probably associated with decreased ocean productivity and prey abundance and therefore decreased breeding capacity.

Nesting season for most populations are constrained temporally (Hamman et al. 2003) and occurs predominantly during the warmer months (Miller 1997). But for some populations, it may occur year-round with seasonal peak and trough periods (Miller 1997). It is generally assumed that nesting season length is shorter at higher latitude beaches (Weishampel et al. 2004) and that nesting activity increases with ocean temperature and photoperiod (Williams-Walls et al. 1983). Nesting seasonality is probably a result of females adjusting their reproductive activity to coincide with conditions that contribute most to maximize reproductive fitness (Cheng 2009 et al.).

The impact of temperature on seasonality of nesting has not yet been thoroughly analysed across species at a global scale. Understanding these impacts will help to better predict the effects of climate change on these species. To address this issue, I reviewed and examined all published information on seasonality across four species of marine turtles: loggerhead (*Caretta caretta*), leatherback (*Dermochelys coriacea*), hawksbill (*Eretmochelys imbricata*) and green turtle (*Chelonia mydas*). The other three species, namely, olive ridley (*Lepidochelys olivacea*), Kemp's ridley (*Lepidochelys kempi*) and flatback turtle (*Natator depressus*) were excluded from this study due to lack of reliable published data from the restricted number of nesting sites under study for these species. The specific question asked was: What are the key drivers of seasonality pattern and duration?

## MATERIALS AND METHODS

**Seasonality of nesting.** A database of the monthly distribution of nests in a season from different regions for four species of marine turtles (*Caretta caretta*, *Chelonia mydas*, *Dermochelys coriacea* and *Eretmochelys imbricata*) was created. The nesting season data were obtained from published literature. Where data were only presented graphically, figures were digitized. Criteria for selecting the data were as follows: 1) data must have covered the full nesting season; 2) only data that provides the number of clutches laid daily, weekly or monthly were included. All data were aggregated by month. In cases where data were for more than one nesting season, the mean number of nests for each month was calculated.

**Duration of season.** An additional database was created for the duration of nesting season (in months). In this case, even data for sites for which only a date range was given could be included.

**Locations.** For locations of all study sites, see Fig. 1-3. Latitude in decimal degrees ca. 5km offshore for each region, were considered as the distance from the equator.

**SST data.** For every site and for the corresponding year-range, SST was obtained from Hadley Ice and Sea Surface Temperature (Hadley ISST) dataset for data prior to 1985 and from Advanced Very High Resolution Radiometer (AVHRR) dataset for data after 1985. For sites for which the year that the nesting season was defined was not given, the mean SST for the five years prior to the publication date was selected.

**Statistical analyses.** Data were analysed using GenStat (11<sup>th</sup> Edition). Test for the assumptions for normal distribution were performed prior to all analyses, and in case assumptions were not met adequate transformations were made. Linear mixed effects models with restricted maximum likelihood (REML) analysis were constructed to determine the factor that dictates the duration of nesting season and the month of peak nesting. Differences were considered significant at  $P < 0.05$ .

## RESULTS

A total of 66 populations from 53 sites (*Caretta caretta* n=17; *Chelonia mydas* n=19; *Dermochelys coriacea* n=19; *Eretmochelys imbricata* n=11) and 115 populations from 97 sites (*C. caretta* n= 20; *C. mydas* n=45; *D. coriacea* n=25; *E. imbricata* n=25) were included for the seasonality of nesting and for the duration of season, respectively. It worth noting that some sites yielded data for more than one species.

### Duration of nest season

The mean season duration in months was  $4.4 \pm 1.2$  (range of 3-8 months) for *Caretta caretta*,  $7.8 \pm 3.4$  (range of 3-12 months) for *Chelonia mydas*,  $6.5 \pm 2.8$  (range of 3-12 months) for *Dermochelys coriacea* and  $7.6 \pm 3.1$  (range of 3-12 months) for *Eretmochelys imbricata* (Fig. 4). Duration was broadly correlated with latitude in all four species (Fig. 5).

To fully elucidate the drivers of seasonal duration I sought to construct a model including species, latitude and mean SST. Due to a high correlation between distance from the equator and mean SST ( $r^2=0.62$ ) the model was first run with both mean SST and distance degrees as fixed effects, however, SST was significant and distance degrees was not but alone both terms were significant. SST and species were incorporated as fixed effects in the linear mixed model, site was included as a random effect. Duration of nesting season was significantly longer with increased mean SST (Linear Mixed Effects Model, Wald statistic = 29.04,  $p < 0.01$ ) but was not significantly affected by species (Linear Mixed Effects Model, Wald statistic = 5.91,  $p = 0.13$ ).

### Seasonality of nesting

In general, nesting seasonality occurred at warmest time of the year (Fig. 6). To elucidate how SST affects seasonality I constructed a model of how the proportion of nesting may have been driven by SST in peak month or each of the 2 months before and after. SST for the month before, month after and the month of peak nesting significantly affected the month of peak nesting (respectively: Linear Mixed Effects Model, Wald statistic = 5.32,  $p < 0.05$ ; Wald statistic = 5.59,  $p < 0.05$ ; and Wald statistic = 6.94,  $p < 0.01$ ). However, the SST two months before and two months after the month of peak nesting did not influence the month of peak nesting.

## DISCUSSION

Temperature and photoperiod will differ in strength and intensity especially along a latitudinal gradient (Hamann et al. 2003). Thus, it is generally assumed that duration of a nesting season (defined as the period of first and the last clutches laid during a year) is determined by latitude, being usually shorter on higher latitudes where summer is comparatively shorter than in the tropics. Although the results of this study showed this was the case, the length of a nesting season was dictated much more by SST: it was generally longer in hotter sites.

However, previous studies reported contrasting findings: Mazaris et al. (2008) found no significant difference between SST and duration of nesting season for the Mediterranean loggerhead population. Pike et al. (2006) found that nesting season was shorter with increasing SST for loggerhead population nesting along the Atlantic coast of central Florida (USA), and Hawkes et al. (2007) reported that the duration of the nesting season in warmer years was significantly longer for loggerheads nesting along Atlantic coast of North Carolina

(USA). Although SST is probably the main predictor of duration of a nesting season, it is possible that there are also other factors, apart from latitude, that determine the duration of a nesting season. This may explain these contrasting findings; however, more research is needed on the inter-relationships of factors affecting duration of a nesting season.

From my research findings, nesting season generally occurred in the warmest months for the four species of marine turtles but there were exceptions. For instance, in Masira Island, Oman (Fig. 2, site 64), the highest proportion of *Caretta caretta* nests occurred in the coolest month. Other exceptions occurred in Mafia Island, Tanzania, where higher proportions of *Chelonia mydas* nests also occurred in relatively colder months (Fig. 2, site 52; Fig. 6-h); and in Playon de Mexiquillo, Mexico, where during the warmest months no nesting occurred for *Dermochelys coriacea* (Fig. 1, site 16; Fig. 6-k).

To paraphrase Owens (1980), “eggs must be on the beach at the right time of year”, because successful incubation is only possible within certain thermal limits (Davenport 1997). Indeed, Godley et al. (2002) reported that for the population of green turtles (*Chelonia mydas*) nesting at Ascension Island, little or no nesting occurs with sand temperatures being close to or below critical limits for development. Furthermore, at warmer temperatures a female is able to re-nest sooner (Hays et al. 2002) as the length of the internesting interval decreases when the water is warmer (Sato et al. 1998). Further research needs to be conducted to look at the possible linkage between the length of internesting interval and the peak nesting month.

Hawkes et al. (2007) suggested that fundamental and realised thermal niches for nesting may not essentially be the same. If this is true, for some populations the optimal time to nest may not be at the warmest month. This also helps to prevent extremely skewed sex ratios. It can be assumed that if the peak of nest season coincides with one of the warmest months of the year, more female offspring will probably be produced. Indeed, Freedberg and Wade (2001) hypothesize that female-biased sex ratios in reptiles with TSD are strengthened by natal nest site fidelity of females. Furthermore, global warming may result in a strong sex ratio bias that could lead to extinction (Hulin et al. 2009). Although shifts in behaviour related to environmental warming are occurring (Weishampel et al. 2004), it remains to be seen if sea turtles could change the timing of reproduction (Hulin et al. 2009).

There are definite advantages to the positive correlation between peak of nesting and warm temperatures. These include better conditions for egg incubation. However, as we have seen, this positive correlation does not always hold true and the reasons for this remain unanswered. Strong evidence has been presented that SST is the key driver of seasonality pattern and duration. Nesting seasons were longer with increased SST. Thus in the context of global warming, it is possible that rookeries that are experiencing an increase in SST may also see an increase in the length of nesting season and strong sex ratio bias. Unfortunately, the current study was limited by the availability of published data. I hope that more data on seasonality across all species of marine turtles will be published. Further research might then investigate the effects of SST on the duration of nest season and on the month of peak nesting for the populations and the species excluded from this analysis.



*Acknowledgements.* This study was supported by University of Exeter funding. I thank M. Witt for assistance with statistics and logistical support. I would also like to thank B.J. Godley for all his support.

## LITERATURE CITED

- Aiken J, Austin T, Broderick A, Godley B, Ebanks-Petrie G (2002) Marine Turtles in the Cayman Islands, British West Indies. In: Mosier A, Foley A, Brost B (eds) Proc 20th Annu Symp Sea Turtle Biol Conserv. NOAA, Florida, p 253-256
- Alkindi A, Mahmoud I, Al-Habsi A, Al-Bahri SN, Al-Gheilani H, Bakheit C (2006) The effect of physical and human factors on beach selection by green turtles (*Chelonia mydas*) at Ras Al-Hadd reserve, Oman. Chelonian Conserv Biol 5:289-294
- Asrar FF (1999) Decline of marine turtle nesting populations in Pakistan. Mar Turtle Newslett 83:13-14
- Bal G, Breheret N, Vanleeuwe H (2007) An update on sea turtle conservation activities in the Republic of Congo. Mar Turtle Newslett 116:9-10
- Baptistotte C, Thomé JC, Bjorndal KA (2003) Reproductive biology and conservation status of the loggerhead sea turtle (*Caretta caretta*) in Espírito Santo State, Brazil. Chel Conserv Biol 4:523-529
- Baran I, Türkozan O (1996) Nesting activity of the loggerhead turtle, *Caretta caretta*, on Fethiye beach, Turkey, in 1994. Chelonian Conserv Biol 2:93-96
- Beasley J, Malpass H, Grant GS (1998) Loggerhead turtle nesting on Topsail Island, North Carolina: 1990-1995. In: Byles R, Fernandez Y (eds) Proc 16th Annu Symp Sea Turtle Biol Conserv. NOAA, Florida, p 16-19
- Beggs JA, Horrocks JA, Krueger BH (2007) Increase in hawksbill sea turtle *Eretmochelys imbricata* nesting in Barbados, West Indies. Endang Species Res 3:159-168
- Bell CD, Solomon JL, Blumenthal JM, Austin TJ, Ebanks-Petrie G, Broderick AC, Godley BJ (2007) Monitoring and conservation of critically reduced marine turtle nesting populations: lessons from the Cayman Islands. Anim Conserv 10:39-47
- Benson SR, Kisokau K, Ambio L, Rei V, Dutton P, Parker D (2007) Beach use, internesting movement, and migration of leatherback turtles, *Dermochelys coriacea*, nesting on the north coast of Papua New Guinea. Chelonian Conserv Biol 6:7-14
- Broderick AC, Coyne MS, Fuller WJ, Glen F, Godley BJ (2007) Fidelity and over-wintering of sea turtles. Proc R Soc Lond B 274:1533–1538
- Broderick Ac, Godley BJ (2000) Temperature and the temporal spread of marine turtle nesting and hatching in Cyprus, Eastern Mediterranean. In: Abreu-Grobis F, Briseño-Dueñas R, Márquez-Millán R, Sarti-Martínez L (eds) Proc 18th Int Sea Turtle Symp. NOAA, Florida, p 181-182
- Bull JJ (1980) Sex determination in reptiles. Q Rev Biol 55:3-21
- Burney CM, Mattison C, Fisberz L (1990) The relationships of loggerhead nesting patterns and moon phase in

- Broward County, Florida. In: Richardson TH, Richardson JL, Donnelly M (eds) Proc 10th Annu Workshop Sea Turtle Biol Conserv. NOAA Technical Memorandum NMFS-SEFC-278, Florida, p 161-164
- Butler JA (2001) Nesting Biology of the sea turtles of St. Kitts, West Indies. *Chelonian Conserv Biol* 4:191-196
- Caldwell DK, (1959) The loggerheads turtles of Cape Romain, South Carolina. In: Austin OL (ed) The Atlantic loggerhead sea turtle, *Caretta caretta* (L.) Bull Fla State Mus Biol Sci 4:319-348
- Carr A, Carr MH (1972) Site fixity in the Caribbean green turtle. *Ecology* 53:425-429
- Catry P, Barbosa C, Indjai B, Almeida A, Godley BJ, Vié J (2002) First census of the green turtle at Poilão, Bijagós Archipelago, Guinea-Bissau: the most important nesting colony on the Atlantic coast of Africa. *Oryx* 36:400-403
- Chacón-Chaverri D, Eckert K (2007) Leatherback sea turtle nesting at Gandoca beach in Caribbean Costa Rica: management recommendations from fifteen years of conservation. *Chelonian Conserv Biol* 6:101-110
- Chaloupka M, Kamezaki N, Limpus C (2008) Is climate change affecting the population dynamics of the endangered Pacific loggerhead sea turtle? *J Exp Mar Biol Ecol* 356:136-143
- Chárriez-Serrano V, Rivera-Muñiz S, Montero-Acevedo L, Moral-Lebrón R (2002) Sea turtle conservation project: hawksbill turtle at Humacao beaches in Puerto Rico. In: Mosier A, Foley A, Brost B (eds) Proc 20th Annu Symp Sea Turtle Biol Conserv. NOAA Technical Memorandum NMFS-SEFSC-477, p 240-243
- Chaves A, Serrano G, Marin G, Arguedas E, Jimenez A, Spotila J (1996) Biology and conservation of leatherback turtles, *Dermochelys coriacea*, at Playa Langosta, Costa Rica. *Chelonian Conserv Biol* 2:184-189
- Cheng IJ, Huang CT, Hung PY, Ke BZ, Kuo CW, Fong CL (2009) Ten years of monitoring the nesting ecology of the green turtle, *Chelonia mydas*, on Lanyu (Orchid Island), Taiwan. *Zool Stud* 48:83-94
- Chu CT, Booth DT, Limpus CJ (2008) Estimating the sex ratio of loggerhead turtle hatchlings at Mon Repos rookery (Australia) from nest temperatures. *Aust J Zool* 56:57-64
- Chua TH (1988) Nesting population and frequency of visits in *Dermochelys coriacea* in Malaysia. *J Herpetol* 22:192-207
- Davenport J (1997) Temperature and the life-history strategies of sea turtles. *J Therm Biol* 22:479-488
- Eckert K, Eckert S (1984) St. Croix leatherback project – 1984 season. *Mar Turtle Newslett* 31:4-6
- Engeman RM, Martin RE, Smith HT, Woolard J, Crady CK, Constantin B, Stahl M, Groninger NP (2006) Impact on predation of sea turtle nests when predator control was removed midway through the nesting season. *Wildl Res* 33:187-192
- Engeman RM, Martin RE, Smith HT, Woolard J, Crady CK, Shwiff SA, Constantin B, Stahl M, Griner J (2005) Dramatic reduction in predation on marine turtle nests through improved predator monitoring and management. *Oryx* 39:318-326
- Freedberg S, Wade MJ (2001) Cultural inheritance as a mechanism for population sex-ratio bias in reptiles. *Evolution* 55:1049-1055
- Girondot M, Rivalan P, Wongsopawiro R, Briane J, Hulin V, Caut S, Guirlet E, Godfrey MH (2006) Phenology of marine turtle nesting revealed by statistical model of the nesting season. *BMC Ecol* 6:11



- Godley BJ, Broderick AC, Frauenstein R, Glen F, Hays GC (2002) Reproductive seasonality and sexual dimorphism in green turtles. *Mar Ecol Prog Ser* 226:125-133
- Hamann M, Limpus CJ, Owens DW (2003) Reproductive cycles of males and females. In: Lutz PL, Musick JA, Wyneken J (eds) *The biology of sea turtles 2*. CRC Press, Boca Raton FL, p 225–241
- Hawkes LA, Broderick AC, Godfrey MH, Godley BJ (2007) Investigating the potential impacts of climate change on a marine turtle population. *Global Change Biol* 13:1-10
- Hays GC, Broderick AC, Glen F, Godley BJ, Houghton JD, Metcalfe JD (2002) Water temperature and interesting intervals for loggerhead (*Caretta caretta*) and green (*Chelonia mydas*) sea turtles. *J Therm Biol* 27: 429–432
- Heppell SS, Snover ML, Crowder LB (2003) Sea turtle population ecology. In: Lutz PL, Musick JA, Wyneken J (eds) *The biology of sea turtles 2*. CRC Press, Boca Raton FL, p 225–241
- Hillis ZM (1990) Buck Island reef national monument sea turtle research program: 1989 – the year of hawksbills and hurricanes. In: Richardson TH, Richardson JI, Donnelly M (eds) *Proc 10th Annu Workshop Sea Turtle Biol Conserv*. NOAA Technical Memorandum NMFS-SEFC-278, p 15-20
- Hilterman ML (2001) *The sea turtles of Suriname, 2000*. Biotopic, Amsterdam, 61 p
- Hirth, HF (1980) Some aspects of the nesting behaviour and reproductive biology of sea turtles. *Am Zool* 20:507-523
- Hirth, HF (1997) Synopsis of the biological data on the green turtle *Chelonia mydas* (Linnaeus 1758). Fish and Wildlife Service, U.S. Department of the Interior, Washington, D. C., 120 p
- Hitipeuw C, Dutton P, Benson S, Thebu J, Bakarbessy J (2007) Population status and interesting movement of leatherback turtles, *Dermochelys coriacea*, nesting on the northwest coast of Papua, Indonesia. *Chelonian Conserv Biol* 6:28-36
- Hughes GR (1974) *The sea turtles of south-east Africa*. Oceanographic Research Institute, Durban, Investigational Report n°36
- Hulin V, Delmas V, Girondot M, Godfrey MH, Guillon JM (2009) Temperature-dependent sex determination and global change: are some species at greater risk? *Oecologia* 160:493–506
- Islam MZ (2006) Sea turtle conservation at St. Martin Island, Bangladesh. In: Pilcher NJ (ed) *Proc 23th Annu Symp Sea Turtle Biol Conservation*. NOAA Technical Memorandum NMFS-536, p 106-108
- Jribi I, Bradai MN, Bouain A (2006) Loggerhead turtle nesting activity in Kuriat Islands (Tunisia): assessment of nine years monitoring. *Mar Turtle Newslett* 112:12-13
- Juarez-Ceron JA, Sarti-Martinez AL, Dutton PH (2003) First study of the green/black turtles of the Revillagigedo Archipelago: a unique nesting stock in the Eastern Pacific. In: Seminoff JA (ed) *Proc 22th Annu Symp Sea Turtle Biol Conserv*. NOAA Technical Memorandum NMFS-SEFSC-503, p 70
- Kamezaki N (1986) Notes on the nesting of the sea turtles in the Yaeyama group, Ryukyu Archipelago. *Jpn J Herpetol* 11:152-155
- Lagueux CJ, Campbell CL, McCoy WA (2003) Nesting and conservation of the hawksbill turtle, *Eretmochelys imbricata* in the Pearl Cays, Nicaragua. *Chelonian Conserv Biol* 4:588-602

- Lazell JD (1980) New England waters: critical habitat for marine turtles. *Copeia* 1980: 290-295
- Limpus CJ (2008) A biological review of Australian marine turtle species 2 Green turtle, *Chelonia mydas* (Linnaeus). Environmental Protection Agency, Queensland, 95 p
- Limpus CJ (2009) A biological review of Australian marine turtle species 3 Hawksbill turtle, *Eretmochelys imbricata* (Linnaeus). Environmental Protection Agency, Queensland, 53 p
- Lopez O, Del Ordi D, Madariaga B, Díaz-Merry A, Ballell L, Abella E, Gracia M, Herraiz L, Borràs S, Varo-Cruz N, Cejudo D, Lopez-Jurado L (2003) Nesting success on the emergences of *Caretta caretta* in the Island of Boavista, Cape Verde, Western Africa. In: Seminoff JA (ed) Sea turtle Symposium 2002. Proc 22th Annu Symp Sea Turtle Biol Conserv. NOAA, Florida, p 150-152
- Lum LL (2005) Beach dynamics and nest distribution of the leatherback turtle (*Dermochelys coriacea*) at Grande Riviere Beach, Trinidad & Tobago. *Rev Biol Trop* 53:239-248
- Marcovaldi MA, Vieitas CF, Godfrey MH (1999) Nesting and conservation management of hawksbill turtles (*Eretmochelys imbricata*) in northern Bahia, Brazil. *Chelonian Conserv Biol* 3:301-307
- Margaritoulis D (2005) Nesting activity and reproductive output of loggerhead sea turtles, *Caretta caretta*, over 19 seasons (1984-2002) at Laganas Bay, Zakynthos, Greece: the largest rookery in the Mediterranean. *Chelonian Conserv Biol* 4:916-929
- Martínez AL, Barragán A, Vargas F, Huerta P, Ocampo E, Tavera A, Escudero A, Vasconcelos D, Ángeles M, Morrison M, Dutton P (2006) The decline of the eastern pacific leatherback and its relation to changes in nesting behaviour and distribution. In: Pilcher NJ (ed) Proc 23th Annu Symp Sea Turtle Biol Conserv. NOAA technical memorandum NMFS-536, p 133- 136
- Mazaris AD, Kallimanis AS, Sgardelis SP, Pantis JD (2008) Do long-term changes in sea surface temperature at the breeding areas affect the breeding dates and reproduction performance of Mediterranean loggerhead turtles? Implications for climate change. *J Exp Mar Biol Ecol* 367:219–226
- McGowan A, Broderick AC, Frett G, Gore S, Hastings M, Pickering A, Wheatley D, White J, Witt MJ, Godley BJ (2008) Down but not out: marine turtles of the British Virgin Islands. *Anim Conserv* 11:92-103
- Miller JD (1997) Reproduction in sea turtles. In: Lutz PL, Musick JA (eds) *Biology of Sea Turtles*. CRC Press, BocaRaton, FL p 51-81
- Miller JD (1985) Embryology of marine turtles. In: Gans C, Billett F, Maderson PFA (eds) *Biology of the Reptilia* Vol 14. John Wiley and Sons, New York, p 270–328
- Milton SL, Lutz PL (2003) Physiological and genetic responses to environmental stress. In: Lutz PL, Musick JA, Wyneken J (eds) *The biology of sea turtles* 2. CRC Press, Boca Raton, FL, p 225–241
- Morreale SJ, Ruiz GJ, Spotila JR, Standora EA (1982) Temperature-dependent sex determination: current practices threaten conservation of sea turtles. *Science*: 216:1245-1247.
- Mortimer JA, Bresson R (1999) Temporal distribution and periodicity in hawksbill turtles (*Eretmochelys imbricata*) nesting at Cousin Island, Republic of Seychelles, 1971-1997. *Chelonian Conserv Biol* 3:318-325
- Mortimer JA, Carr A (1987) Reproduction and Migrations of the Ascension Island Green Turtle (*Chelonia mydas*). *Copeia* 1987:103-113
- Mrosovsky N (1980) Thermal Biology of Sea Turtles. *Am Zool* 20:531-547

- Mrosovsky N, Hopkins-Murphy SR, Richardson JI (1984) Sex ratio of sea turtles: seasonal changes. *Science* 225:739-741
- Mrosovsky N, Dutton PH, Whitmore CP (1984b) Sex ratios of two species of sea turtle nesting in Suriname. *Can J Zool* 62:2227-2239
- Muir C, Abdallah O (2006) Community-based conservation of sea turtles at Mafia Island, Tanzania. In: Pilcher NJ (compiler) *Proc 23th Annu Symp Sea Turtle Biol Conserv*. NOAA Technical Memorandum NMFS-536, p 79-81
- Niethammer KR, Balazs GH, Hatfield JS, Nakai GL, Megyesi J (1997) Reproductive biology of the green turtle (*Chelonia mydas*) at Tern Island, French Frigate Shoals, Hawai'i. *Pac Sci* 51:36-47
- Ordoñez C, Troëng S, Meylan A, Meylan P, Ruiz A (2007) Chiriqui beach, Panama, the most important leatherback nesting beach in central America. *Chelonian Conserv Biol* 6:122-126
- Owens D (1980) The comparative reproductive physiology of sea turtles. *Am Zool* 20:549-563
- Peñate JG, Karamoko M, Bamba S, Djadji G (2007) An update on marine turtles in Côte d'Ivoire, West Africa. *Mar Turtle Newslett* 11:7-8
- Perez-Castañeda R, Salum-Fares A, Defeo O (2007) Reproductive patterns of the hawksbill turtle *Eretmochelys imbricata* in sandy beaches of the Yucatan Peninsula. *J Mar Biol Ass U K* 87:815-824
- Pike DA, Stiner JC (2007) Sea turtle species vary in their susceptibility to tropical cyclones. *Oecologia* 153:471-478
- Pike DA, Antworth RL, Stiner JC (2006) Earlier nesting contributes to shorter nesting seasons for the loggerhead seaturtle, *Caretta caretta*. *J Herpetol* 40:91-94
- Pilcher NJ (1999) The hawksbill turtle, *Eretmochelys imbricata*, in the Arabian Gulf. *Chelonian Conserv Biol* 3:312-317
- Pilcher NJ, Ali L (1999) Reproductive biology of the hawksbill turtle, *Eretmochelys imbricata*, in Sabah, Malaysia. *Chelonian Conserv Biol* 3:330-336
- Plotkin PT (2003) Adult migrations and habitat use. In: Lutz PL, Musick JA, Wyneken J (eds) *The biology of sea turtles 2*. CRC Press, Boca Raton, FL, p 225-241
- Rees AF, Saad A, Jony M (2008) Discovery of a regionally important green turtle *Chelonia mydas* rookery in Syria. *Oryx* 42:456-459
- Rivera-Muñiz S, Montero-Acevedo L, Charriez-Serrano V, Moral-Lebron R (2002) Leatherback sea turtle (*Dermochelys coriacea*) at Humacao beaches in Puerto Rico. In: Mosier A, Foley A, Brost B (eds). *Proc 20th Annu Symp Sea Turtle Biol Conserv*. NOAA Technical Memorandum NMFS-SEFSC-477, p 237-240
- Ross JP (1979) Sea turtles in the Sultanate of Oman. *WWF Project 1320*, 53p
- Sato K, Bando T, Matsuzawa Y, Tanaka H, Sakamoto W, Minamikawa S, Goto K (1997) Decline of the loggerhead turtle, *Caretta caretta*, nesting on Senri Beach in Minabe, Wakayama, Japan. *Chelonian Conserv Biol* 2:600-603
- Sato K, Matsuzawa Y, Tanaka H, Bando T, Minamikawa S, Sakamoto W, Naito Y (1998) Internesting intervals for loggerhead turtles, *Caretta caretta*, and green turtles, *Chelonia mydas*, are affected by temperature. *Can J Zool* 76:1651-1662

- Scott JA (2006) Use of Satellite telemetry to determine ecology and management of loggerhead turtle (*Caretta caretta*) during the nesting season in Georgia. MSc dissertation, University of Georgia, Athens, GA
- Solow AR, Bjorndal KA, Bolten AB (2002) Annual variation in nesting numbers of marine turtles: the effect of sea surface temperature on re-migration intervals. *Ecol Lett* 5: 742–746
- Standora EA, Spotila JR (1895) Temperature dependent sex determination in sea turtles. *Copeia* 1985:711-722
- Suganuma H, Kamezaki N, Yusuf A (1999) Current status of nesting populations of the hawksbill turtle (*Eretmochelys imbricata*). *Chelonian Conserv Biol* 3:337-343
- Sunderraj SF, Joshua J (2006) Status report on the UNEP – CMS sea turtle project on the Gujarat coast, India. *Indian Ocean Turtle Newslett* 3:6-10
- Thomé J, Baptistotte C, Moreira L, Scalfoni J, Almeida A, Rieth D, Barata P (2007) Nesting biology and conservation of the leatherback sea turtle (*Dermochelys coriacea*) in the State of Espírito Santo, Brazil, 1988–1989 to 2003–2004. *Chelonian Conserv Biol* 6:15-27
- Tiwari M, Bjorndal KA, Bolten AB, Bolker BM (2006) Evaluation of density-dependent processes and green turtle *Chelonia mydas* hatchling production at Tortuguero, Costa Rica. *Mar Ecol Prog Ser* 326:283-293
- Tomás J, Castroviejo J, Raga JA (1999) Sea turtles in the south of Bioko Island (Equatorial Guinea). *Mar Turtle Newslett* 84:4-6
- Weishampel JF, Bagley DA, Ehrhart LM (2004) Earlier nesting by loggerhead sea turtles following sea surface warming. *Global Change Biol* 10:1424-1427
- Williams-Walls N, O'Hara J, Gallagher RM, Worth DF, Peery BD, Wilcox JR (1983) Spatial and temporal trends of sea turtle nesting of Hutchinson Island, Florida, 1971-1979. *Bull Mar Sci* 33:55-66
- Witt MJ, Baert B, Broderick AC, Formia A, Fretey J, Gibudi A, Moussounda C, Mounguengui G, Nguouessomo S, Pamell R, Roumet D, Sounguet G, Verhage B, Zogo A, Godley BJ (in press) Aerial surveying of the world's largest leatherback turtle rookery: a more effective methodology for large-scale monitoring. *Biol Conserv*
- Witzell WN (1983) Synopsis of biological data on the hawksbill turtle *Eretmochelys imbricata* (Linnaeus, 1766). *FAO Fish Synop* (137):78 p
- Yalçın-Özdilek Ş (2007) Status of sea turtles (*Chelonia mydas* and *Caretta caretta*) on Samandağ beach, Turkey: a five-year monitoring study. *Ann Zool Fennici* 44:333-347
- Yasuda T, Tanaka H, Kittiwattanawong K, Mitamura H, Klom-in W, Arai N (2006) Do female green turtles (*Chelonia mydas*) exhibit reproductive seasonality in a year-round nesting rookery? *J Zool* 269:451-457
- Yntema CL, Mrosovsky N (1982) Critical periods and pivotal temperatures for sexual differentiation in loggerhead sea turtles. *Can J Zool* 60:1012-1016
- Zambrano R, Rodríguez E (1995) The nesting biology of hawksbill (*Eretmochelys imbricata*) and green (*Chelonia mydas*) sea turtles in El Cuyo, Yucatan, Mexico. In: Richardson JI, Richardson TH (eds) *Proc 12<sup>th</sup> Annu Workshop Sea Turtle Biol Conserv*. NOAA Technical Memorandum NMFS-SRFSC-361, p 252-256

## FIGURE LEGENDS

Fig. 1. Nesting sites in the Americas. *Caretta caretta*: 1=Topsail Island, USA (Beasley et al. 1998) 2=Cape Romain, U.S.A (Caldwell et al. 1959) 3=Georgia, USA (Scott 2006) 4=Canaveral National Seashore, Florida, USA (Pike & Stiner 2007) 5=Jupiter Island, USA (Engeman et al. 2006) 6= Broward County, Florida (Burney et al. 1990) 10=Cayman Islands (Aiken et al. 2002) 26= Buritaca, Colombia (Hirth 1980) 42=Espírito Santo State, Brazil (Baptistotte et al. 2003). *Chelonia mydas*: 4=Canaveral National Seashore, Florida, U.S.A (Pike & Stiner 2007) 5= Jupiter Island, Florida, USA (Engeman et al. 2005) 8= El Cuyo, Yucatan, Mexico (Zambrano & Rodríguez 1995) 9= Cuba (Hirth 1997) 10= Cayman Islands (Bell et al. 2007) 11= Dominican Republic (Hirth 1997) 12=Puerto Rico (Hirth 1997) 15=Revillagigedo Archipelago, Mexico (Juarez-Ceron et al. 2003) 19=St. Kitts, West Indies (Butler 2001) 20=Belize (Hirth 1997) 21=Martinique (Hirth 1997) 23=El Salvador (Hirth 1997) 29=Tortuguero, Costa Rica (Tiwari et al. 2006) 35=Krofajapasi, Suriname (Mrosovsky et al. 1984b) 38=Colombia (Hirth 1997) 39=Atol das Rocas, Brazil (Hirth 1997) 41=Praia do Forte, Brazil (Hirth 1997). *Dermochelys coriacea*: 4=Canaveral National Seashore, Florida, USA (Pike & Stiner 2007) 13= Humacao Beaches, Puerto Rico (Rivera-Muñiz et al. 2002) 14= British Virgin Islands (McGowan et al. 2008) 16= Playon de Mexiquillo, Mexico (Martinez et al. 2006) 18= Sandy Point, U.S. Virgin Islands (Eckert & Eckert 1984) 19= St. Kitts, West Indies (Butler 2001) 27= Trinidad Island (Hirth 1980) 28=Grande Riviere Beach, Trinidad and Tobago (Lum 2005) 30=Playa Langosta, Costa Rica (Chaves et al. 1996) 32=Gandoca Beach, Costa Rica (Chacón-Chaverri & Eckert 2007) 33=Chiriqui Beach, Panama (Ordoñez et al. 2007) 36=Baboensanti, Suriname (Hilterman 2001) 37=Ya:lima:po beach, French Guiana (Girondot et al. 2006) 43= State of Espírito Santo (north coast), Brazil (Thomé et al. 2007). *Eretmochelys imbricata*: 7= The Bahamas (Witzell 1983) 8= El Cuyo, Yucatan, Mexico (Pérez-Castañeda et al. 2007) 13= Humacao Beaches, Puerto Rico (Chárriez-Serrano et al. 2002) 17= Buck Island Reef National Monument, U.S.A (Hillis 1990) 19= St. Kitts, West Indies (Butler 2001) 22= Saint Lucia (Witzell 1983) 24=Barbados (Beggs et al. 2007) 25=Pearl Cays, Nicaragua (Lagueux et al. 2003) 31=Venezuela (Witzell 1983) 34=Suriname (Witzell 1983) 38=Colombia (Witzell 1983) 40=Arembepe and Praia do Forte, Bahia, Brazil (Marcovaldi et al. 1999). Solid circles indicate sites for which data were available on seasonality and season duration. Open circles indicate sites for which only seasonal duration were available.

Fig. 2. Nesting sites in Africa and Middle East. *Caretta caretta*: 44=Boavista island, Cape Verde (Lopez et al 2003) 54=Tongaland, South Africa (Hughes 1974) 55= Kuriat Islands, Tunisia (Jribi et al. 2006) 56= Laganas Bay, Zakynthos island, Greece (Margaritoulis, D. 2005) 57= Fethiye Beach, Turkey (Baran and Türkozan 1996) 58= Samandağ Beach, Turkey (Yalçın-Özdilek 2007) 60= Alagadi Beach, Cyprus (Broderick & Godley 2000) 64= Masira, Oman (Ross 1979). *Chelonia mydas*: 45=Poilão, Bijagós Archipelago, Guinea-Bissau (Catry et al. 2002) 46=Ivory Coast (Peñate et al. 2007) 47=South of Bioko Island, Equatorial Guinea (Tomás et al. 1999) 49=Republic of Congo (Bal et al. 2007) 50=Ascension Island (Mortimer & Carr 1987) 52=Mafia Island, Tanzania (Muir & Abdallah 2006) 53=Aldabra Atoll (Hirth 1997) 58=Samandağ Beach, Turkey (Yalçın-Özdilek 2007) 59=Latakia beach, Siria (Rees et

al. 2008) 60=Alagadi Beach, Cyprus (Broderick & Godley 2000) 63=Saih Al-Marai, Oman (Alkindi et al. 2006). *Dermochelys coriacea*: 46=Ivory Coast (Peñate et al. 2007) 47=South of Bioko Island, Equatorial Guinea (Tomás et al. 1999) 48=Gamba, Gabon (Witt et al. in press) 49=Republic of Congo (Bal et al. 2007) 54=Tongaland, South Africa (Hughes 1974). *Eretmochelys imbricata*: 47=South of Bioko Island, Equatorial Guinea (Tomás et al. 1999) 51=Cousin Island, Seychelles (Mortimer & Bresson 1999) 61= Karan, Kurayn, Jana and Jurayd islands, Arabian Gulf (Pilcher 1999) 65= Oman (Witzell 1983) 66=Jabal Aziz, South Yemen (Hirth 1980). Solid circles indicate sites for which data were available on seasonality and season duration. Open circles indicate sites for which only seasonal duration were available.

Fig. 3. Nesting sites in Asia and Oceania. *Caretta caretta*: 70= Ryukyu Archipelago (Kamezaki 1986) 74= Senri Beach, Japan (Sato et al. 1997) 95= Mon Repos, Australia (Chu et al. 2008). *Chelonia mydas*: 62=Hawkes Bay and Sandspit, Pakistan (Asrar 1999) 67=Gujarat coast, India (Sunderraj & Joshua 2006) 68= St. Martin Island, Bangladesh (Islam 2006) 69= Lanyu (Orchid Island), Taiwan (Cheng et al. 2009) 71= Xisha Islands, China (Hirth 1997) 75=Tern Island, Hawaii (Niethammer et al. 1997) 77=Huyong Island, Thailand (Yasuda et al. 2006) 78=Sri Lanka, Kosgoda (Hirth 1997) 81=Sarawak, Malaysia (Hirth 1997) 82=Maldives(Hirth 1997) 86=Phoenix Islands (Hirth 1997) 87=Long Island, Papua New Guinea (Hirth 1997) 89=Berau Islands, Indonesia (Hirth 1997) 92= Gulf of Carpentaria, Australia (Limpus 2008) 93= Northern Great Barrier Reef, Australia (Limpus 2008) 94=South Queensland, Australia (Limpus 2008) 97=Fiji (Hirth 1997). *Dermochelys coriacea*: 80=Rantau Abang, Malaysia (Chua 1988) 83=Jamursba-Medi, Papua, Indonesia (Hitipeuw et al. 2007) 84=Wermon, Papua, Indonesia (Hitipeuw et al. 2007) 88=Kamiali WMA, Papua New Guinea (Benson et al. 2007) 96=Southeastern Queensland, Australia (Limpus 2009). *Eretmochelys imbricata*: 70= Ryukyu Archipelago (Kamezaki 1986) 72= Philippines (Witzell 1983) 73=Thailand (Witzell 1983) 76= Palau Is, Micronesia (Hirth 1980) 79=Pulau Gulisaan, Malaysia (Pilcher & Ali 1999) 85=Gresik, Indonesia (Suganuma et al. 1999) 90=Solomon Islands, Australia (Limpus 2009) 91=Torres Strait, Australia (Limpus 2009). Solid circles indicate sites for which data were available on seasonality and season duration. Open circles indicate sites for which only seasonal duration were available. For logistical purposes site 86 (Phoenix Islands) is not shown.

Fig. 4. Range of temperature for nesting (a) and for the month of peak nesting (b) for each species. *Caretta caretta* (Cc), *Chelonia mydas* (Cm), *Dermochelys coriacea* (Dc) and *Eretmochelys imbricata* (Ei).

Fig. 5. Relationships between latitude degrees from the equator and duration of nest season (transformed to normalize data) for each species. (a) *Caretta caretta* ( $r^2 = 0.13$ ), (b) *Chelonia mydas* ( $r^2 = 0.27$ ), (c) *Dermochelys coriacea* ( $r^2 = 0.29$ ) and (d) *Eretmochelys imbricata* ( $r^2 = 0.03$ ).



Fig. 6. Seasonal distribution of nests and variation of SST. *Caretta caretta*: a – Jupiter Island, Florida (U.S.A); b – Grand Cayman, Cayman Islands; c – Espírito Santo state, Brazil; d – Boavista Island, Cape Verde; e – Alagadi beach, Northern Cyprus. *Chelonia mydas*: f – Jupiter Island, Florida (U.S.A); g – Tortuguero, Costa Rica; h – Mafia Island, Tanzania; i – Huyong Island, Thailand; j – Tern Island, Hawaii. *Dermochelys coriacea*: k – Playon de Mexiquillo, Mexico; l – St. Kitts, West Indies; m – Gandoca Beach, Costa Rica; n – Gamba, Gabon; o – South of Bioko Island, Equatorial Guinea. *Eretmochelys imbricata*: p – El Cuyo, Yucatan Peninsula, Mexico; q – Barbados; r – Pearl Cays, Nicaragua; s – Arembepe and Praia do Forte, Bahia, Brazil; t – Arabian Gulf (islands: Karan, Kurayn, Jana and Jurayd).

## FIGURES




Fig. 1




Fig. 2




Fig. 3





Fig.4




Fig.5






Fig. 6